

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 7/50, A61K 38/12	A1	(11) International Publication Number: WO 97/05163 (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/US96/12111 (22) International Filing Date: 23 July 1996 (23.07.96) (30) Priority Data: 08/506,790 25 July 1995 (25.07.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/506,790 (CON) Filed on 25 July 1995 (25.07.95) (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indi- anapolis, IN 46285 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): RODRIGUEZ, Michael, J. [US/US]; 1825 Sailing Court, Indianapolis, IN 46260 (US). (74) Agents: MCCLAIN, Janet, T. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: CYCLIC PEPTIDE ANTI-FUNGAL AGENTS (57) Abstract Provided are pharmaceutical formulations, and methods of inhibiting fungal and parasitic activity using a compound of formula (I) or a pharmaceutically acceptable salt thereof.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

CYCLIC PEPTIDE ANTI-FUNGAL AGENTS

This invention relates to semi-synthetic cyclic peptide compounds which are useful as antifungal agents and which have improved stability and water solubility. In particular, it relates to derivatives of the echinocandin class of cyclic peptides; to methods for treating fungal infections, and to formulations useful in the methods.


The compounds provided by this invention are semi-synthetic compounds derived from cyclic peptides which are produced by culturing various microorganisms. A number of cyclic peptides are known in the art including echinocandin B (A30912A), aculeacin, mulundocandin, sporiofungin and S31794/F1.

In general, these cyclic peptides may be structurally characterized as a cyclic hexapeptide core (or nucleus) with an acylated amino group on one of the core amino acids. The amino group is typically acylated with a fatty acid group forming a side chain off the nucleus. For example, echinocandin B has a linoleoyl side chain while aculeacin has a palmitoyl side chain.

The fatty acid side chains may be removed from the cyclic peptide core to provide an amino nucleus (for example, a compound of formula I, below, where R^2 is hydrogen). The amino group may then be re-acylated to provide semi-synthetic compounds such as those claimed in the present application.

The echinocandin B nucleus has been re-acylated with certain non-naturally occurring side chain moieties to provide a number of antifungal agents (see, Debono, U.S. Pat. Ser. No. 4,293,489). Among such antifungal agents is cilofungin which is represented by a compound of formula I where R^{z1} and R^{z2} are each $-CHOHCH_3$, R^{z3} is methyl; R^{x1} is hydrogen, R^{x2} , R^{y1} , R^{y2} , R^{y3} , R^{y4} and R^0 is hydroxy and R_2 is p-(octyloxy)benzoyl.

The present invention provides a compound of formula I:



5 wherein:

R^{z1} is hydrogen, $-CH_2OH$, $-CHOHCH_3$ or $-CHOHCH_2C(O)NH_2$;

R^{z2} is hydrogen, $-CH_2OH$ or $-CHOHCH_3$;

R^{z3} is hydrogen or methyl;

R^{x1} is hydrogen, hydroxy or $O-R^{x1'}$;

10 $R^{x1'}$ is C_1-C_6 alkyl, benzyl, $-(CH_2)_2Si(CH_3)_3$,
 $-CH_2CH=CH_2$, $-CH_2CHOHCH_2OH$, $-(CH_2)_aCOOH$, $-(CH_2)_bNR^{w1}R^{w2}$,
 $-(CH_2)_cPOR^{w3}R^{w4}$ or $-[(CH_2)_2O]_d-(C_1-C_6)alkyl$;

a , b and c are independently 1, 2, 3, 4, 5 or 6;

R^{w1} and R^{w2} are independently hydrogen, C_1-C_6 alkyl,

15 or R^{w1} and R^{w2} combine to form $-CH_2(CH_2)_eCH_2-$;


R^{w3} and R^{w4} are independently hydroxy, or C_1-C_6
 alkoxy;

d is 1 or 2;


e is 1, 2 or 3;

20 R^{x2} , R^{y1} , R^{y2} , R^{y3} and R^{y4} are independently hydroxy or
 hydrogen;





R^0 is hydroxy, $-OP(O)(OH)_2$ or a group of the formulae:



R^1 is C_1 - C_6 alkyl, phenyl, p-halo-phenyl, p-nitrophenyl, benzyl, p-halo-benzyl or p-nitro-benzyl;



R^3 is



R^{3a} , R^{3b} , R^{3c} and R^{3d} are independently hydrogen, C_1 - C_{12} alkyl, C_2 - C_{12} alkynyl, C_1 - C_{12} alkoxy, C_1 - C_{12} alkylthio, halo, $-\text{O}-(\text{CH}_2)_m-[\text{O}-(\text{CH}_2)_n]_p-\text{O}-(\text{C}_1\text{-C}_{12} \text{ alkyl})$ or $-\text{O}-(\text{CH}_2)_q-\text{X}-\text{R}^4$;

m is 2, 3 or 4;

n is 2, 3 or 4;

p is 0 or 1;

q is 2, 3 or 4;

X is pyrrolidino, piperidino or piperazino; and

R^4 is hydrogen, C_1 - C_{12} alkyl, C_3 - C_{12} cycloalkyl, benzyl or C_3 - C_{12} cycloalkylmethyl;

with the proviso that at least one of R^{z1} and R^{z2} must be hydrogen;

or a pharmaceutically acceptable salt thereof.

Also provided are pharmaceutical formulations, methods for inhibiting fungal activity and methods of treating fungal infections which employ the compounds of the invention.

As used herein, the term "C₁-C₁₂ alkyl" refers to a straight or branched alkyl chain having from one to twelve carbon atoms. Typical C₁-C₁₂ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, t-butyl, 5 pentyl, 5-methylpentyl, hexyl, heptyl, 3,3-dimethylheptyl, octyl, 2-methyl-octyl, nonyl, decyl, undecyl, dodecyl and the like. The term "C₁-C₁₂ alkyl" includes within its definition the term "C₁-C₆ alkyl."

The term "halo" refers to chloro, fluoro, bromo or 10 iodo.

The term "C₂-C₁₂ alkynyl" refers to a straight or branched alkynyl chain having from two to twelve carbon atoms. Typical C₂-C₁₂ alkynyl groups include ethynyl, 1-propyn-1-yl, 1-propyn-2-yl, 1-butyn-1-yl, 1-butyn-3-yl, 1- 15 pentyn-3-yl, 4-pentyn-2-yl, 1-hexyn-3-yl, 3-hexyn-1-yl, 5-methyl-3-hexyn-1-yl, 5-octyn-1-yl, 7-octyn-1-yl, 4-decyn-1-yl, 6-decyn-1-yl and the like.

The term "C₁-C₁₂ alkylthio" refers to a straight or branched alkyl chain having from one to twelve carbon atoms 20 attached to a sulfur atom. Typical C₁-C₁₂ alkylthio groups include methylthio, ethylthio, propylthio, isopropylthio, butylthio, 3-methyl-heptylthio, octylthio, 5,5-dimethyl-hexylthio and the like.

The term "C₁-C₁₂ alkoxy" refers to a straight or 25 branched alkyl chain having from one to twelve carbon atoms attached to an oxygen atom. Typical C₁-C₁₂ alkoxy groups include methoxy, ethoxy, propoxy, butoxy, sec-butoxy, pentoxy, 5-methyl-hexoxy, heptoxy, octyloxy, decyloxy dodecyloxy and the like. The term "C₁-C₁₂ alkyl" includes 30 within its definition the term "C₁-C₆ alkoxy."

The term "C₃-C₁₂ cycloalkyl" refers to a saturated hydrocarbon ring structure having from three to twelve carbon atoms. Typical C₃-C₁₂ cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 35 cycloheptyl, cyclooctyl and the like. The term "C₃-C₁₂ cycloalkylmethyl" refers to a C₃-C₁₂ cycloalkyl attached to a methylene group.

The term "amino-protecting group" as used in the specification refers to substituents of the amino group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. Examples of such amino-protecting groups include formyl, trityl, phthalimido, trichloroacetyl, chloroacetyl, bromoacetyl, iodoacetyl groups, or urethane-type blocking groups such as benzyloxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, t-butoxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(p-toluy)-prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)-ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)-ethoxycarbonyl, fluorenylmethoxycarbonyl ("Fmoc"), 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl, 1-piperidyloxycarbonyl and the like; benzoylmethylsulfonyl, 2-nitrophenylsulfenyl, diphenylphosphine oxide and like amino-protecting groups. The species of amino-protecting group employed is not critical so long as the derivatized amino group is stable to the condition of subsequent reaction(s) on other positions of the intermediate molecule and can be selectively removed at the appropriate point without disrupting the remainder of the molecule including

any other amino-protecting group(s). Preferred amino-protecting groups are *t*-butoxycarbonyl (*t*-Boc), allyloxycarbonyl and benzyloxycarbonyl (CbZ). Further examples of groups referred to by the above terms are
5 described by J. W. Barton, "Protective Groups in Organic Chemistry", J. G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, Chapter 2, and T. W. Greene, "Protective Groups in Organic Synthesis", John Wiley and sons, New York, N.Y., 1981, Chapter 7.

10 The term "hydroxy protecting group" refers to a substituent of an hydroxy group that is commonly employed to block or protect the hydroxy functionality while reactions are carried out on other functional groups on the compound. Examples of such hydroxy protecting groups
15 include tetrahydropyranyl, 2-methoxyprop-2-yl, 1-ethoxyethyl-yl, methoxymethyl, β -methoxyethoxymethyl, methylthiomethyl, *t*-butyl, *t*-amyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, 4,4',4"-trimethoxytrityl, benzyl, allyl, trimethylsilyl, trimethylsilylethyl, (*t*-
20 butyl)dimethylsilyl, and 2,2,2-trichloroethoxycarbonyl and the like. The species of hydroxy protecting group is not critical so long as the derivatized hydroxy group is stable to the conditions of the subsequent reaction(s) and can be removed at the appropriate point without disrupting the
25 remainder of the molecule. A preferred hydroxy protecting group is trimethylsilylethyl. Further examples of hydroxy protecting groups are described in T.W. Greene, "Protective Groups in Organic Synthesis," John Wiley and Sons, New York, N.Y., (2nd ed., 1991) chapters 2 and 3. The term
30 "protected hydroxy" refers to a hydroxy group bonded to one of the above hydroxy protecting groups.

The term "inhibiting", i.e. a method of inhibiting fungal activity, includes stopping, retarding or prophylactically hindering or preventing the growth or any
35 attending characteristics and results from the existence of a fungus.

The term "contacting", i.e. contacting a compound of the invention with a fungus, includes a union or junction, or apparent touching or mutual tangency of a compound of the invention with a fungus. However, the term does not
5 imply any further limitations to the process, such as by mechanism of inhibition, and the methods are defined to encompass the spirit of the invention, which is to inhibit fungal activity by the action of the compounds and their inherent antifungal properties, or in other words, the
10 compounds, used in the claimed methods are the causative agent for such inhibition.

The term "pharmaceutically acceptable salt" as used herein, refers to salts of the compounds of the above formula which are substantially non-toxic to living
15 organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a mineral or organic acid or an inorganic base. Such salts are known as acid addition and base addition salts.

20 Acids commonly employed to form acid addition salts are mineral acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid and the like, and organic acids such as *p*-toluenesulfonic, methanesulfonic acid, oxalic acid, *p*-bromophenylsulfonic
25 acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate,
30 pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate,
35 chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, ph nylacetate, phenylpropionate,

phenylbutyrate, citrate, lactate, γ hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like. Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic acids such as maleic acid and methanesulfonic acid.

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like. The potassium and sodium salt forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is not of a critical nature, so long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

Preferred compounds of this invention are those compounds of formula I where:

R^{z1} is hydrogen, $-CH_2OH$ or $-CHOHCH_3$;

or a pharmaceutically acceptable salt thereof.

Of these compounds, more preferred are those compounds of formula I where:

R^{z1} and R^{z2} are independently hydrogen or $-CHOHCH_3$;

R^{z3} is methyl;

R^{y1} , R^{y2} , R^{y3} and R^{y4} are each hydroxy;

R^{x1} is hydrogen, hydroxy or $-OR^{x1'}$;


$R^{x1'}$ is methyl, benzyl, $-CH_2CHOHCH_2OH$, $-CH_2COOH$,

$-(CH_2)_2NR^{w1}R^{w2}$ or $-(CH_2)_2POR^{w3}R^{w4}$;

R^{w1} and R^{w2} are independently hydrogen or methyl;

R^{w3} and R^{w4} are independently hydrogen or methoxy;

R^0 is hydroxy or a group of the formulae:



R^1 is methyl;

5 or a pharmaceutically acceptable salt thereof.


Of these preferred compounds, further preferred are those compounds of formula I where:

R^{z1} and R^{z2} are both hydrogen;

R^{x1} is hydrogen;

10 R^{x2} is hydrogen;

R^3 is





R^{3c} and R^{3d} are independently hydrogen, C_1 - C_{12} alkyl, C_2 - C_{12} alkynyl, C_1 - C_{12} alkoxy or

15 $\text{---O---(CH}_2\text{)}_2\text{---O---(C}_1\text{---C}_{12}\text{ alkyl)}$;

or a pharmaceutically acceptable salt thereof.

Of these compounds the most preferred compounds are those compounds where:

R^3 is




20

or a pharmaceutically acceptable salt thereof.


The compounds of formula I may be prepared according to Reaction Scheme I, as follows.

25

Reaction Scheme I

(IA)

A. deacylate




(IB)


5

B. re-acylate

11



C. Retro-aldol
reaction



wherein:

- 5 R^{nat} is a naturally occurring cyclic peptide sidechain;

R' is hydrogen, methyl or $-CH_2C(O)NH_2$;

R'' is hydrogen or methyl; and

R^{z1}, R^{z2}, R^{z3}, R^{x1}, R^{x1'}, R^{x2}, R^{y1}, R^{y2}, R^{y3}, R^{y4}, R⁰ and R² are as defined above.

Reaction scheme I, above, is accomplished by carrying out reactions A-C, above. Once a reaction is complete, the intermediate compound may be isolated by procedures well-known in the art, for example, the compound may be crystallized or precipitated and then collected by filtration, or the reaction solvent may be removed by extraction, evaporation or decantation. The intermediate compound may be further purified, if desired, by common techniques such as crystallization or precipitation or chromatography over solid supports such as silica gel, alumina and the like, before carrying out the next step of the reaction scheme.

In reaction IA, a naturally occurring cyclic peptide of the formula IA is deacylated using procedures known in the art to provide an amino nucleus of formula IB. This reaction is typically carried out using enzymatic deacylation by exposing the naturally occurring cyclic peptide to a deacylase enzyme. The deacylase enzyme may be obtained from the microorganism *Actinoplanes utahensis* and used substantially as described in U.S. Patent Nos. 4,293,482 and 4,304,716, herein incorporated by reference. The deacylase enzyme may also be obtained from the *Pseudomonas* species. Deacylation may be accomplished using whole cells of *Actinoplanes utahensis* or *Pseudomonas* or the crude or purified enzyme thereof or using an immobilized form of the enzyme. See European Patent Application No. 0 460 882 (December 11, 1991). Examples of naturally occurring cyclic peptides which may be used as starting materials include aculeacin (palmitoyl side chain), tetrahydroechinocandin B (stearoyl side chain), mulundocandin (branched C₁₅ side chain), L-671,329 (C₁₆ branched side chain), S 31794/F1 (tetradecanoyl side chain), sporiofungin (C₁₅ branched side chain), FR901379 (palmitoyl side chain) and the like. A preferred naturally occurring cyclic peptide is chinocandin B (a compound of

formula IA where R', R" and R^{z3} are each methyl, R^{x1}, R^{x2}, R^{y1}, R^{y2}, R^{y3}, R^{y4} and R⁰ are each hydroxy and R² is linoleoyl).

5 In Reaction IB, the amino nucleus of formula IB is re-acylated using procedures known in the art to provide a compound of formula IC where R² is an acyl group as defined hereinabove.

For example, the amino nucleus may be acylated by reaction with an appropriately substituted acyl halide, preferably in the presence of an acid scavenger such as a tertiary amine, such as triethylamine. The reaction is typically carried out at a temperature of from about -20°C to about 25°C. Typical solvents for this reaction include polar aprotic solvents such as dioxane or
10 dimethylformamide. Solvent choice is not critical so long as the solvent employed is inert to the ongoing reaction and the reactants are sufficiently solubilized to effect the desired reaction.

The amino nucleus may also be acylated by reaction
20 with an appropriately substituted carboxylic acid, in the presence of a coupling agent. Typical coupling agents include dicyclohexylcarbodiimide (DCC), N,N'-carbonyldiimidazole, bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP-Cl), N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), benzotriazol-1-yloxytripyrrolidinophosphonium
25 hexafluorophosphate (PyBOP) and the like.


In addition, the amino nucleus may be acylated with an activated ester of a carboxylic acid such as an ester of a carboxylic acid of the formula R²-COOH and p-nitrophenyl,
30 2,4,5-trichlorophenyl, hydroxybenzotriazole hydrate (HOBT·H₂O), pentafluorophenol, N-hydroxysuccinimide and the like. Preferred acylating moieties are the active esters of the carboxylic acid R²-COOH such as 2,4,5-trichlorophenyl ester and benzotriazole ester. The
35 reaction is typically carried out for one to sixty five hours at a temperature from about 0°C to about 30°C in an aprotic solvent. The reaction is generally complete after

about twenty four to forty eight hours when carried out a temperature of from about 15°C to about 30°C. Typical solvents for this reaction are tetrahydrofuran and dimethylformamide or a mixture of such solvents. The amino
5 nucleus is generally employed in equimolar proportions relative to the activated ester or with a slight excess of the amino nucleus.

In Reaction IC, one or both of the β -hydroxy groups are selectively removed from a compound of formula IC to
10 provide a compound of formula I. Specifically, the compound of formula IC is heated in an aprotic solvent to between about 50°C and 100°C in the presence of at least about five equivalents (preferably at least ten
15 equivalents) of a retro-aldol-promoting reagent selected from the group consisting of trimethylamine-N-oxide, triethylamine-N-oxide, trimethylamine-N-oxide-hydrate and trimethylamine-N-oxide-hydrate. The reaction is generally conducted in a sealed tube or in a round bottom flask supplied with a water condenser for approximately three to
20 seventy two hours. The reaction is preferably conducted at a temperature of about 70°C to about 100°C, preferably at about 70°C. Typical aprotic solvents include dimethylformamide, tetrahydrofuran, acetonitrile, dimethylsulfoxide and the like. A preferred solvent is
25 acetonitrile. Preferred retro-aldol-promoting reagents are trimethylamine-N-oxide and trimethylamine-N-oxide-hydrate. The most preferred retro-aldol-promoting reagent is trimethylamine-N-oxide-hydrate. This reaction is preferably conducted after protecting the amination hydroxy
30 (R^{x1}) of the compound of formula IC. The hydroxy protecting group may be removed after removal of the β -hydroxy group(s), using procedures known in the art.

The term "aldol" indicates a molecule that is both an alcohol and an aldehyde or ketone wherein the hydroxyl and
35 carbonyl functional groups are on adjacent carbon atoms as shown below. Thus, as used herein, the term " β hydroxy group" refers to the alcohol moiety including both the

β -carbon and the hydroxyl functional group that is resident on the β -carbon according to the following general structure:



5

where:

R^1 and R^2 are independently hydrogen or methyl;

R^3 and R^4 taken together with the atoms to which they are attached form a cyclic peptide.

- 10 The compounds where R^{x1} is $-O-R^{x1'}$ may be prepared by the corresponding compound where R^{x1} is hydroxy with an appropriately substituted alcohol in the presence of an acid to provide a compound of formula I wherein R^{x1} is $-O-R^{x1'}$ where $R^{x1'}$ is C_1-C_6 alkyl, benzyl, $-(CH_2)_2Si(CH_3)_3$,
 15 $-CH_2CH=CH_2$, $-(CH_2)_aCOOH$, $-(CH_2)_bNR^{w1}R^{w2}$, $-(CH_2)_cPOR^{w3}R^{w4}$ or $-[(CH_2)_2O]_d-(C_1-C_6)alkyl$. The compounds where R^{x1} is $-O-R^{x1'}$ are preferably prepared after removal of the β -hydroxy group(s). The reaction is typically carried out in a polar aprotic solvent such as dioxane or
 20 dimethylsulfoxide at a temperature of from about $0^\circ C$ to about $35^\circ C$, preferably at about room temperature. Solvent choice is not critical so long as the solvent employed is inert to the ongoing reaction and the reactants are sufficiently solubilized to effect the desired reaction.
 25 Preferred acids include p-toluenesulfonic acid, hydrochloric acid and camphorsulfonic acid.

The compounds where $R^{x1'}$ is $-(CH_2)_bNR^{w1}R^{w2}$ where R^{w1} and R^{w2} are hydrogen may be prepared via a protected compound wherein R^{x1} is $-(CH_2)_bNHR^a$ where R^a is an amino

protecting group. The resultant protected compound may be deprotected according to procedures known in the art.

The compounds where $R^{x1'}$ is $-\text{CH}_2\text{CHOHCH}_2\text{OH}$ may be prepared by hydroxylating a compound of formula I where
5 $R^{x1'}$ is $-\text{CH}_2\text{CH}=\text{CH}_2$ with osmium tetroxide in the presence of a catalyst at a temperature in the range of from about 0°C to about 40°C for about one to twenty four hours in a organic/aqueous solvent mixture, for example dioxane/water. Suitable catalysts include N-methylmorpholine N-oxide (NMO)
10 and the like. Typical solvents suitable for use in this reaction include dimethylformamide, tetrahydrofuran, ethyl acetate, dioxane or a mixture of these solvents. Solvent choice is not critical so long as the solvent employed is inert to the ongoing reaction and the reactants are
15 sufficiently solubilized to effect the desired reaction. The reaction is preferably conducted at a temperature in the range of from about 20°C to about 30°C for about eighteen to twenty four hours.

The compounds where R^0 is hydroxy may be
20 phosphorylated by reaction with an appropriately substituted alkyl, phenyl or benzyl phosphate to provide a compound where R^0 is $-\text{O}-\text{P}(\text{O})\text{OH}-\text{OR}^1$ where R^1 is C_1-C_6 alkyl, phenyl or benzyl, or by reaction with an appropriately substituted alkyl, phenyl or benzyl phosphonic acid to
25 provide a compound where R^0 is $-\text{O}-\text{P}(\text{O})\text{OH}-\text{R}^1$ where R^1 is C_1-C_6 alkyl, phenyl or benzyl, to provide a compound of formula I where R^0 is a group of the formula $-\text{OP}(\text{O})\text{OH}-\text{OR}^1$ or $-\text{OP}(\text{O})\text{OH}-\text{R}^1$. This reaction is preferably conducted after removal of the β -hydroxy group(s). The phosphonic
30 acid is typically used in an activated form, for example as a phosphonic halide, preferably a phosphonic chloride. The reaction is carried out in the presence of a base such as lithium trimethylsilanolate (LiOTMS), lithium bis(trimethylsilyl)amide (LHMDS), pyridine and the like.
35 The reaction is typically carried out for up to one hour at a temperature from about -30°C to about 0°C in an aprotic

solvent such as tetrahydrofuran and dimethylformamide. The reaction is generally complete in about fifteen minutes when carried out under these conditions. The phosphate or phosphonate reactant is generally employed in equimolar proportions to about a one mole excess relative to the amino nucleus in the presence of an equimolar or slight excess of the base. Phosphorylation of an amino nucleus with unprotected aminal hydroxy groups is typically carried out at lower temperatures, for example from about -30°C to about -15°C.

Alternatively, the compounds where R^{X1} is hydroxy may be phosphorylated after first protecting the hydroxy moiety with an hydroxy protecting group using procedures known in the art. For example, the reaction is typically carried out by combining the compound with a suitable hydroxy protecting group in the presence of a catalyst at a temperature in the range of from about 0°C to about 40°C for about one to five hours in a mutual inert solvent. The hydroxy protecting group is generally employed in an amount ranging from about equimolar proportions to about a 100 molar excess relative to the compound, preferably in a large molar excess. Suitable catalysts include strong acids such as p-toluenesulfonic acid, camphorsulfonic acid (CSA), hydrochloric acid, sulfuric acid, trifluoroacetic acid and the like. Typical solvents suitable for use in this reaction include any organic solvent such as dioxane. Solvent choice is not critical so long as the solvent employed is inert to the ongoing reaction and the reactants are sufficiently solubilized to effect the desired reaction. The reaction is preferably conducted at a temperature in the range of from about 20°C to about 30°C for about two to four hours. The protected compound is then phosphorylated as described above. The hydroxy protecting group is then removed according to procedures known in the art to provide the desired phosphorylated compound. For example, the protecting groups can be removed by reaction with a Lewis acid in a mutual inert

organic solvent such as methylene chloride. Examples of Lewis acids include trimethylsilylbromide, boron trifluoride etherate and the like. The reaction is typically carried out at a temperature of from about 0°C to about 40°C, preferably at a temperature of from about 20°C to about 30°C. A preferred Lewis acid is boron trifluoride etherate.

The dideoxy compounds are prepared by removing the benzylic and aminor hydroxy groups (R^{x2} and R^{x1} , respectively). The hydroxy groups may be removed by subjecting a non-dideoxy compound to a strong acid and a reducing agent at a temperature of between -5°C and 70°C, in a suitable solvent. Typical strong acids include trichloroacetic acid, trifluoroacetic acid or borontrifluoride etherate. A preferred strong acid is trifluoroacetic acid. Typical reducing agents include sodium cyanoborohydride or triethylsilane. A preferred reducing agent is triethylsilane. Suitable solvents include methylene chloride, chloroform or acetic acid, preferably methylene chloride. The strong acid should be present in an amount of from 2 to 80 mol per mol of substrate, and the reducing agent should be present in an amount of 2 to 80 mol per mol of substrate. This process affords selective removal of the aminor and benzylic hydroxy groups.

The cyclic peptides used to make the compounds of the present invention may be prepared by fermentation of known microorganisms. For example, the cyclic peptide of formula IB where R' , R'' and R^{z3} are methyl, and R^{x1} , R^{x2} , R^{y1} , R^{y2} , R^{y3} , R^{y4} and R^0 are each hydroxy (cyclic nucleus corresponding to A-30912A) may be prepared using the procedure detailed in Abbott et al., U.S. Pat. Ser. No. 4,293,482, which is herein incorporated by reference. The cyclic peptide of formula IB where R' , R'' and R^{z3} are methyl, R^{x1} is hydroxy, R^{x2} is hydrogen, and R^{y1} , R^{y2} , R^{y3} , R^{y4} and R^0 are each hydroxy (cyclic nucleus corresponding to A-30912B) may be prepared using the procedure detailed


in Abbott et al., U.S. Pat. Ser. No. 4,299,763, which is herein incorporated by reference. Aculeacin may be prepared using the procedure detailed in Mizuno et al., U.S. Pat. Ser. No. 3,978,210 which is herein incorporated
5 by reference. The cyclic peptide of formula IB where R' is -CH₂C(O)NH₂, R'' is methyl, R^{z3} is hydrogen, and R^{x1}, R^{x2}, R^{y1}, R^{y2}, R^{y3}, R^{y4} and R⁰ are each hydroxy may be prepared by deacylating the cyclic peptide prepared using the procedure detailed in Chen et al., U.S. Pat. Ser. No.
10 5,198,421, which is herein incorporated by reference.

The R²-COOH precursor acids, used to acylate the amino nucleus of formula IB, may be obtained commercially or prepared according to procedures known in the art. For example, an appropriately substituted phenyl boronic acid
15 or biphenyl boronic acid reactant may be reacted with a p-halobenzoic acid reactant in the presence of a catalyst such as tetrakis(triphenylphosphine)palladium and an inorganic base such as potassium carbonate in a mutual inert organic solvent such as toluene at a temperature of
20 from about 20°C to the reflux temperature of the reaction mixture to provide the corresponding biphenyl carboxylic acids and terphenyl carboxylic acids used to prepare the compounds of formula I. The reaction is typically carried out with equimolar proportions of the boronic acid reactant
25 and the p-benzoic acid reactant, or a slight molar excess of the p-benzoic acid reactant relative to the boronic acid reactant, and a 1-2 molar excess of the inorganic base. The reaction is generally complete after about four to about ten hours when carried out at reflux temperature in
30 toluene.

The boronic acid reactant may be prepared by reacting an appropriately substituted halophenyl or halobiphenyl reactant with two equivalents of triisopropyl borate in the presence of an alkyl lithium, for example sec-butyl
35 lithium, in a mutual inert solvent such as tetrahydrofuran. The alkyl lithium is typically employed in a slight molar excess relative to the halophenyl or halobiphenyl reactant.



The alkyl lithium is typically combined with the solvent by dropwise addition at reduced temperatures ($<-70^{\circ}\text{C}$) and allowed to stir for approximately thirty minutes before the addition of the triisopropyl borate. The reaction is typically carried out initially at a temperature of from about -100°C to about -50°C , preferably from about -75°C to about -85°C for thirty minutes to two hours and then warmed to room temperature and reacted for an additional one to three hours. The reaction is generally complete in from several minutes to about four hours. When the reaction is substantially complete, the boronic acid moiety is formed by the addition of an acid. A preferred acid is a 1N hydrochloric acid solution.

The carboxylic acids, $\text{R}^2\text{-COOH}$, having an acetylene moiety may be prepared by reacting an appropriately substituted acetylene reactant with an appropriately substituted phenyl or biphenyl reactant of the formula



where L is a suitable leaving group such as bromo, iodo, methanesulfonate, toluenesulfonate, trifluoromethanesulfonate and the like, in the presence of a catalyst and preferably in the presence of an acid scavenger in a mutual inert solvent such as acetonitrile. Examples of acid scavengers include triethylamine and pyridine, preferably triethylamine. A preferred catalyst is formed *in situ* from palladium (II) chloride, triphenylphosphine and copper (I) iodide. The reaction is typically carried out for thirty minutes to twenty one hours at a temperature from about room temperature to the reflux temperature of reaction mixture. The reaction is generally complete after about two to about six hours when carried out at reflux temperature.

Alternatively, a suitably substituted phenyl reactant of the formula $\text{halo}-\text{C}_6\text{H}_4-\text{L}$ may be reacted with an appropriately substituted acetylene reactant as described above to provide, for example, a compound of the formula


halo--C≡C--R^{3c} which can be coupled with a phenyl boronic acid reactant as described above.

The following Preparations and Examples further describe how to synthesize the compounds of the present invention. The terms melting point, proton nuclear magnetic resonance spectra, mass spectra, infrared spectra, ultraviolet spectra, elemental analysis, high performance liquid chromatography, and thin layer chromatography are abbreviated "m.p.", "NMR", "MS", "IR", "UV", "Analysis", "HPLC", and "TLC", respectively. In addition, the absorption maxima listed for the IR spectra are only those of interest and not all of the maxima observed.

Example 1

15

A. Preparation of the compound of formula IC where R', R" and R²³ are each methyl, R^{x1} and R^{x2} are each hydroxy, R^{y1}, R^{y2}, R^{y3} and R^{y4} are each hydroxy, R⁰ is hydroxy and R² is




20 The N-acyl cyclic peptide derivatives listed in Table 3, below were prepared by dissolving 348.1 g (60.2 mmol) of the A30912A nucleus (compound of formula IB where R', R" and R²³ are each methyl, R^{x1} and R^{x2} are each hydroxy, R^{y1}, R^{y2}, R^{y3} and R^{y4} are each hydroxy, and R⁰ is hydroxy) and 26.0 g
25 (48.2 mmol) of the 2,4,5-trichlorophenol ester of [(4"-pentyloxy)-1,1':4',1"-terphenyl]-4-carboxylic acid, in 8.5 l of dimethylformamide. The resultant reaction mixture was stirred for approximately 48 hours at room temperature and then the solvent was removed in vacuo to provide a residue.
30 This residue was slurried in ether, collected by filtration, washed with methylene chloride and then dissolved in methanol or a 1:1 (v/v) acetonitrile/water mixture. The resultant solution is subjected to reverse phase HPLC (C18; eluent of 20-40% aqueous acetonitrile
35 containing 0.5% monobasic ammonium phosphate (w/v);

20 ml/min.; 230 nm). After removing the unreacted A30912A nucleus, the desired product was eluted from the column using an eluent of aqueous acetonitrile. The fractions containing the desired product are combined and then
 5 reduced to dryness in vacuo to provide the desired acylated nucleus.

Yield: 18 g.

MS(FAB): 1140.5103 (M⁺).

- 10 B. Preparation of the compound of formula IC where R', R" and R^{z3} are each methyl, R^{x1} and R^{x2} are each hydrogen, R^{y1}, R^{y2}, R^{y3} and R^{y4} are each hydroxy, R^{z1} is hydroxy and R^{z2} is




- To a mixture of 5 g (4.4 mmol) of the compound
 15 prepared in Example 1A and 17 ml of trifluoroacetic acid in 250 ml of methylene chloride, was added 35 ml of triethylsilane. When the reaction was substantially complete, as indicated by HPLC (C18, eluent of 55% acetonitrile; 2 ml/min; 280 nm; R_T (starting material) =
 20 4.19 min.; R_T (product) = 6.40 min.), the reaction mixture was concentrated in vacuo to provide a solid. This solid was slurried in 100 ml of 50% aqueous acetone and then dissolved by adjusting the pH of the mixture to
 25 approximately pH 7. The resultant solution was poured into a large volume of water (approximately 1 liter) resulting in the precipitation of a white solid. This solid was isolated by filtration through a sintered glass funnel, washed with diethyl ether and then dried in vacuo at 55°C to provide 3.718 g of the titled compound. The funnel was
 30 washed with methanol to collect the remaining solid, which was dried in vacuo to provide an additional 0.154 g of the titled compound.

Yield: 3.872 g (79%).

MS (FAB): m/e 1108.7 (M)

- 35 HPLC: (eluent of 55% acetonitrile; 2 ml/min.; 280 nm):
 R_T = 6.43 min.


- 5 C. Preparation of the compound of formula I where R^{z1} and R^{z2} are each hydrogen, R^{z3} is methyl, R^{x1} and R^{x2} are each hydrogen, R^{y1}, R^{y2}, R^{y3} and R^{y4} are each hydroxy, R⁰ is hydroxy and R² is



- 10 The compound prepared in Example 1B was dissolved in a 1:1 mixture of acetonitrile and dimethylformamide. To this solution, was added TNO-hydrate (1.6 g, 14.3 mmol) all at once. The reaction mixture was heated at 100°C for 48 hours after which the mixture was cooled to room temperature and concentrated to approximately one-half its original volume. The crude residue was dissolved with acetic acid and purified by reverse phase preparative HPLC to yield 150 mg final product (52%). TNO-hydrate converted the two threonine residues into glycine residues and left the homotyrosine untouched as confirmed by fast atom bombardment mass spectrometry, which confirmed the molecular weight of the product having the chemical formula
- 20 C₅₄H₆₅N₇O₁₃: calculated - 1020.1; found - 1020.8. (M+H).

Example 2

- 25 Alternative Preparation of the compound of formula I where R^{z1} and R^{z2} are each hydrogen, R^{z3} is methyl, R^{x1} and R^{x2} are each hydrogen, R^{y1}, R^{y2}, R^{y3} and R^{y4} are each hydroxy, R⁰ is hydroxy and R² is



- 30 The compound prepared in Example 1B was dissolved in 20 ml of a 1:1 mixture of acetonitrile and dimethylformamide. To this solution, was added trimethylamine N-oxide dihydrate (2.51 g, 22.6 mmol) all at once. The reaction mixture was heated at 100°C for 24 hours in a sealed tube after which the mixture was cooled to room temperature and concentrated in vacuo to approximately one-half its original volume. The crude
- 35

residue was isolated by filtration and then rinsed with cold dimethylformamide. The resultant filtrate was dried in vacuo to provide a residue. This residue was purified using reverse phase HPLC (3x40x10 radial column, eluent of 50% acetonitrile in water, 60 ml/min., 280 nm) to yield 330 mg final product (72%).

Fast atom bombardment mass spectrometry confirmed the molecular weight of the product having the chemical formula $C_{54}H_{65}N_7O_{13}$: calculated - 1020.1; found - 1020.4 (M+H).

10

The compounds of formula I exhibit antifungal activity. For example, the compounds of formula I inhibit the growth of various infectious fungi including Candida spp. such as *C. albicans*, *C. parapsilosis*, *C. krusei*, *C. glabrata*, or *C. tropicalis*, *C. lusitaniae*; *Torulopus* spp. such as *T. glabrata*; *Aspergillus* spp. such as *A. fumigatus*; *Histoplasma* spp. such as *H. capsulatum*; *Cryptococcus* spp. such as *C. neoformans*; *Blastomyces* spp. such as *B. dermatitidis*; *Fusarium* spp., *Trichophyton* spp., *Pseudallescheria boydii*, *Coccidioides immitis*, *Sporothrix schenckii* and the like.

Antifungal activity of a test compound is determined in vitro by obtaining the minimum inhibitory concentration (MIC) of the compound using a standard agar dilution test or a disc-diffusion test. The compound is then tested in vivo (in mice) to determine the effective dose of the test compound for controlling a systemic fungal infection.

Accordingly, the compound prepared in Example 1C was tested for antifungal activity against *C. albicans*. The minimal inhibitory concentration (MIC) of this compound against *C. albicans* was 0.312 (µg/ml).

In addition, the effective dose of the compound prepared in Example 1C, for controlling a systemic fungal infection (*C. albicans*), was tested in vivo (mice). The ED₅₀ for this compound was >2.5.

The compounds of formula I are active in vitro and in vivo and are useful in combating either systemic fungal

infections or fungal skin infections. Accordingly, the present invention provides a method of inhibiting fungal activity comprising contacting a compound of formula I, or a pharmaceutically acceptable salt thereof, with a fungus.

5 A preferred method includes inhibiting *Candida albicans* or *Aspergillus fumigatis* activity. The present invention further provides a method of treating a fungal infection which comprises administering an effective amount of a compound of formula I, or a pharmaceutically acceptable

10 salt thereof, to a host in need of such treatment. A preferred method includes treating a *Candida albicans* or *Aspergillus fumigatis* infection.

With respect to antifungal activity, the term "effective amount," means an amount of a compound of the present invention which is capable of inhibiting fungal activity. The dose administered will vary depending on

15 such factors as the nature and severity of the infection, the age and general health of the host and the tolerance of the host to the antifungal agent. The particular dose regimen likewise may vary according to such factors and may be given in a single daily dose or in multiple doses during the day. The regimen may last from about 2-3 days to about 2-3 weeks or longer. A typical daily dose (administered in

20 single or divided doses) will contain a dosage level of from about 0.01 mg/kg to about 100 mg/kg of body weight of an active compound of this invention. Preferred daily doses generally will be from about 0.1 mg/kg to about 60 mg/kg and ideally from about 2.5 mg/kg to about 40 mg/kg.

The present invention also provides pharmaceutical formulations useful for administering the antifungal compounds of the invention. Accordingly, the present invention also provides a pharmaceutical formulation comprising one or more pharmaceutically acceptable carriers, diluents or excipients and a compound of claim 1.

30 The active ingredient in such formulations comprises from 0.1% to 99.9% by weight of the formulation, more generally from about 10% to about 30% by weight. By

35

"pharmaceutically acceptable" it is meant that the carrier, diluent or excipient is compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

5 A compound of formula I may be administered parenterally, for example using intramuscular, sub-cutaneous, or intra-peritoneal injection, nasal or oral means. In addition to these methods of administration, a compound of formula I may be applied topically for skin
10 infections.

For parenteral administration the formulation comprises a compound of formula I and a physiologically acceptable diluent such as deionized water, physiological saline, 5% dextrose and other commonly used diluents. The
15 formulation may contain a solubilizing agent such as a polyethylene glycol or polypropylene glycol or other known solubilizing agent. Such formulations may be made up in sterile vials containing the antifungal and excipient in a dry powder or lyophilized powder form. Prior to use, a
20 physiologically acceptable diluent is added and the solution withdrawn via syringe for administration to the patient.

The present pharmaceutical formulations are prepared by known procedures using known and readily available
25 ingredients. In making the compositions of the present invention, the active ingredient will generally be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a
30 diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups,
35 aerosols, (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories,

sterile injectable solutions, sterile packaged powders and the like.

For oral administration, the antifungal compound is filled into gelatin capsules or formed into tablets. Such tablets may also contain a binding agent, a dispersant or other suitable excipients suitable for preparing a proper size tablet for the dosage and particular antifungal compound of the formula I. For pediatric or geriatric use the antifungal compound may be formulated into a flavored liquid suspension, solution or emulsion. A preferred oral formulation is linoleic acid, cremophor RH-60 and water and preferably in the amount (by volume) of 8% linoleic acid, 5% cremophor RH-60, 87% sterile water and a compound of formula I in an amount of from about 2.5 to about 40 mg/ml.

For topical use the antifungal compound may be formulated with a dry powder for application to the skin surface or it may be formulated in a liquid formulation comprising a solubilizing aqueous liquid or non-aqueous liquid, e.g., an alcohol or glycol.

The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way. The term "active ingredient" means a compound according to formula I or a pharmaceutically acceptable salt thereof.

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
Active ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

Formulation 2

A tablet is prepared using the ingredients below:

Quantity

	<u>(mg/capsule)</u>
Active ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
5 Stearic acid	<u>5</u>
Total	665 mg

The components are blended and compressed to form tablets each weighing 665 mg.

10 Formulation 3

An aerosol solution is prepared containing the following components:

	<u>Weight</u>
Active ingredient	0.25
15 Methanol	25.75
Propellant 22 (Chlorodifluoromethane)	<u>74.00</u>
Total	100.00

20 The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

25

Formulation 4

Tablets, each containing 60 mg of active ingredient, are made as follows:

Active ingredient	60 mg
30 Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
35 Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
Total	150 mg

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinyl-pyrrolidone is mixed with the resultant powder, and the mixture then is passed
5 through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S.
10 sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

Formulation 5

Capsules, each containing 80 mg of active ingredient,
15 are made as follows:

Active ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
20 Total	200 mg

The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

25

Formulation 6

Suppositories, each containing 225 mg of active ingredient, are made as follows:

Active ingredient	225 mg
30 Saturated fatty acid glycerides	<u>2,000 mg</u>
Total	2,225 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat
35 necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

Formulation 7

Suspensions, each containing 50 mg of active ingredient per 5 ml dose, are made as follows:

5	Active ingredient	50 mg
	Sodium carboxymethyl cellulose	50 mg
	Syrup	1.25 ml
	Benzoic acid solution	0.10 ml
	Flavor	q.v.
10	Color	q.v.
	Purified water to total	5 ml

The active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Formulation 8


20 An intravenous formulation may be prepared as follows:

Active ingredient	100 mg
Isotonic saline	1,000 ml

25 The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 ml per minute.

CLAIMS

1. A compound of formula I:



I

5

wherein:

R^{z1} is hydrogen, $-CH_2OH$, $-CHOHCH_3$ or $-CHOHCH_2C(O)NH_2$;

R^{z2} is hydrogen, $-CH_2OH$ or $-CHOHCH_3$;

R^{z3} is hydrogen or methyl;

10 R^{x1} is hydrogen, hydroxy or $O-R^{x1'}$;

$R^{x1'}$ is C_1 - C_6 alkyl, benzyl, $-(CH_2)_2Si(CH_3)_3$,

$-CH_2CH=CH_2$, $-CH_2CHOHCH_2OH$, $-(CH_2)_aCOOH$, $-(CH_2)_bNR^{w1}R^{w2}$,

$-(CH_2)_cPOR^{w3}R^{w4}$ or $-[(CH_2)_2O]_d-(C_1-C_6)$ alkyl;

a , b and c are independently 1, 2, 3, 4, 5 or 6;

15 R^{w1} and R^{w2} are independently hydrogen, C_1 - C_6 alkyl,

or R^{w1} and R^{w2} combine to form $-CH_2(CH_2)_eCH_2-$;


R^{w3} and R^{w4} are independently hydroxy, or C_1 - C_6 alkoxy;

d is 1 or 2;


20 e is 1, 2 or 3;

R^{x2} , R^{y1} , R^{y2} , R^{y3} and R^{y4} are independently hydroxy or hydrogen;





R^0 is hydroxy, $-OP(O)(OH)_2$ or a group of the formulae:



R^1 is C_1 - C_6 alkyl, phenyl, p-halo-phenyl, p-nitrophenyl, benzyl, p-halo-benzyl or p-nitro-benzyl;

R^2 is ---C(O)---  ---R^3 ;

5 R^3 is



R^{3a} , R^{3b} , R^{3c} and R^{3d} are independently hydrogen, C_1 - C_{12} alkyl, C_2 - C_{12} alkynyl, C_1 - C_{12} alkoxy, C_1 - C_{12} alkylthio, halo, $\text{---O---(CH}_2\text{)}_m\text{---[O---(CH}_2\text{)}_n\text{]}_p\text{---O---(C}_1\text{---C}_{12}\text{ alkyl)}$ or

10 $\text{---O---(CH}_2\text{)}_q\text{---X---R}^4$;

m is 2, 3 or 4;

n is 2, 3 or 4;

p is 0 or 1;

q is 2, 3 or 4;

15 X is pyrrolidino, piperidino or piperazino; and

R^4 is hydrogen, C_1 - C_{12} alkyl, C_3 - C_{12} cycloalkyl, benzyl or C_3 - C_{12} cycloalkylmethyl;

with the proviso that at least one of R^{z1} and R^{z2} must be hydrogen;


20 or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 where:

R^{z1} is hydrogen, $\text{---CH}_2\text{OH}$ or ---CHOHCH_3 ;



or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 2 where:
 R^{z1} and R^{z2} are independently hydrogen or $-\text{CHOHCH}_3$;
 R^{z3} is methyl;
 R^{y1} , R^{y2} , R^{y3} and R^{y4} are each hydroxy;
 5 R^{x1} is hydrogen, hydroxy or $-\text{OR}^{x1'}$;
 $R^{x1'}$ is methyl, benzyl, $-\text{CH}_2\text{CHOHCH}_2\text{OH}$, $-\text{CH}_2\text{COOH}$,
 $-(\text{CH}_2)_2\text{NR}^{w1}\text{R}^{w2}$ or $-(\text{CH}_2)_2\text{POR}^{w3}\text{R}^{w4}$;
 R^{w1} and R^{w2} are independently hydrogen or methyl;
 R^{w3} and R^{w4} are independently hydrogen or methoxy;
 10 R^0 is hydroxy or a group of the formulae:




- R^1 is methyl;
 or a pharmaceutically acceptable salt thereof.
 15

4. A compound according to claim 3 where:
 R^{z1} and R^{z2} are both hydrogen;
 R^{x1} is hydrogen;
 R^{x2} is hydrogen;
 20 R^3 is



- R^{3c} and R^{3d} are independently hydrogen, $\text{C}_1\text{-C}_{12}$ alkyl,
 $\text{C}_2\text{-C}_{12}$ alkynyl, $\text{C}_1\text{-C}_{12}$ alkoxy or $-\text{O}-(\text{CH}_2)_2-\text{O}-(\text{C}_1\text{-C}_{12} \text{ alkyl})$;
 or a pharmaceutically acceptable salt thereof.
 25

5. A compound according to claim 4 where:
 R^3 is



or a pharmaceutically acceptable salt thereof.

5

6. A pharmaceutical formulation comprising a compound of formula I, or a pharmaceutically acceptable salt thereof, as claimed in any one of claims 1 to 5, associated with one or more pharmaceutically acceptable carriers, diluents or excipients therefor.


10

7. A compound of formula I, or a pharmaceutically acceptable salt thereof, as claimed in any one of claims 1 to 5, for use as a pharmaceutical.

15

8. A process for preparing a compound of formula I, or a pharmaceutically acceptable salt thereof, as claimed in any one of claims 1 to 5, comprising heating a compound of formula IC:

20



wherein:

R', R'', RX¹, RX², RY¹, RY², RY³, RY⁴ and R⁰ are as defined in claim 1;
in an aprotic solvent to between 50°C and 100°C in the presence of at least five equivalents of a retro-aldol-promoting reagent selected from the group consisting of trimethylamine-N-oxide, triethylamine-N-oxide, trimethylamine-N-oxide-hydrate and trimethylamine-N-oxide-hydrate;
and then optionally converting the compound of formula I to a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/12111

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 7/50; A61K 38/12

US CL : 514/9, 11; 530/317

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/9, 11; 530/317

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE, STN

search terms: anti-pneumocystis, cyclohexapeptide

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	WO 96/15142 A1 (ELI LILLY AND COMPANY) 23 May 1996 (23.05.96), see entire document, especially page 13.	1-8
A	WO 94/25045 A1 (MERCK & CO., INC.) 10 November 1994 (10.11.94), see entire document.	1-7
A	US 5,166,135 A (SCHMATZ) 24 November 1992 (24.11.92), see entire document.	1-7
A	EP 0,459,564 A2 (MERCK & CO., INC.) 04 December 1991 (04.12.91), see entire document.	1-8
A	EP 0,561,639 A1 (ELI LILLY AND COMPANY) 22 September 1993 (22.09.93), see entire document.	1-8

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

02 OCTOBER 1996

Date of mailing of the international search report

29 OCT 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ANISH GUPTA

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/12111

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94/25050 A1 (MERCK & CO., INC.) 10 November 1994 (10.11.94), see entire document.	1-8